

MATHEMATISCHES FORSCHUNGSINSTITUT OBERWOLFACH

Report No. 25/2005

Nonlinear Evolution Problems

Organised by
Klaus Ecker (Berlin)
Jalal Shatah (New York)
Michael Struwe (Zürich)

May 29th – June 4th, 2005

ABSTRACT. In this workshop three types of nonlinear evolution problems—geometric evolution equations (essentially of parabolic type), nonlinear hyperbolic equations, and dispersive equations— were the subject of 22 talks.

Mathematics Subject Classification (2000): 47J35, 74J30, 35L70, 53C44, 35Q55, 35-06.

Introduction by the Organisers

The workshop ‘Nonlinear Evolution Problems’ focussed on three types of nonlinear evolution equations:

- (1) Geometric evolution equations (essentially of parabolic type)
- (2) Nonlinear hyperbolic equations
- (3) Dispersive equations

The programme consisted of 22 talks presented by international specialists from Australia, France, Germany, Italy, Sweden, Switzerland and the United States. Generally, three lectures were delivered in the morning sessions and two in the late afternoon which left ample time for individual discussions.

In the first group of equations, in particular the Ricci flow and conformal flows such as the Yamabe flow and the Q-curvature flow were considered. A further focus was on curvature flows for hypersurfaces such as the Gauss and harmonic mean curvature flow and on geometric flows of higher order. Julie Clutterbuck presented a direct approach to certain fundamental estimates for a general class of parabolic equations. Mete Soner pointed out important relations between fully nonlinear parabolic equations and backward stochastic differential equations.

The class of hyperbolic equations was represented by wave maps, Einstein's equations of gravitation and nonlinear wave equations in waveguides and cones.

Among the dispersive equations the nonlinear Schroedinger equation and the KdV equation were considered.

The individual discussions covered connections between approaches to the different equations in each group but also on common techniques used across the three classes of equations. Among such techniques are for instance the careful examination of the algebraic and geometric structure of the nonlinear terms. Another common theme appears to be the phenomenon of blow-up (singularity formation) for solutions and methods to describe the solution near these. Selfsimilar solutions and blow-up rates play an important role here.

The organizers decided to particularly encourage the young researchers among the participants to present their work, including a PhD student and several recent post-doctoral fellows.